



**Physiological and environmental controls on the nitrogen and
oxygen isotope fractionation of nitrate during its assimilation by
marine phytoplankton**

by
Kristen Leigh Karsh
Institute for Marine and Antarctic Studies

Submitted in fulfilment of the requirements for the Doctor of Philosophy
University of Tasmania July 2014

Under a conjoint arrangement with Princeton University

Declaration of Originality

This thesis is submitted to both the University of Tasmania and Princeton University as part of a co-tutelle agreement. To the best of my knowledge and belief no material previously published or written by another person except where due acknowledgement is made in the text of the thesis, nor does the thesis contain any material that infringes copyright.

Signed _____ Date _____

Authority of Access

This thesis may be made available for loan and limited copying and communication in accordance with the Copyright Act 1968.

Signed _____ Date _____

Ethical Conduct

The research associated with this thesis abides by the international and Australian codes on human and animal experimentation, the guidelines by the Australian Government's Office of the Gene Technology Regulator, and the rulings of the Safety, Ethics, and Institutional Biosafety Committees of the University.

Signed _____ Date _____

Published Work

The publishers of the papers comprising Chapters 2 and 3 hold the copyright for that content, and access to the material should be sought from the respective journals. The remaining non-published content of the thesis may be made available for loan and limited copying and communication in accordance with the Copyright Act 1968.

Signed _____ Date _____

Abstract

Physiological and environmental controls on the nitrogen and oxygen isotope fractionation of nitrate during its assimilation by marine phytoplankton

by

Kristen Leigh Karsh

Advisers: Thomas Trull and Daniel Sigman

Nitrogen (N) is an essential nutrient for phytoplankton growth and its availability often limits primary production in the surface ocean. Quantifying the inputs, losses, and internal cycling of N is thus a major goal in marine biogeochemical research. The N isotope effect ($^{15}\epsilon_{\text{org}}$) for nitrate assimilation is a key parameter in using N isotope distributions ($^{15}\text{N}/^{14}\text{N}$) to study the marine N cycle. Unexplained variability in its magnitude is a major source of uncertainty to N isotopic studies in the modern and past ocean. The ratio of the oxygen (O) and N isotope effects for nitrate assimilation ($^{18}\epsilon_{\text{org}}/^{15}\epsilon_{\text{org}}$) is also an important parameter; the association of nitrate assimilation with an $^{18}\epsilon_{\text{org}}/^{15}\epsilon_{\text{org}}$ near 1 is the cornerstone of studies using coupled nitrate N and O isotope measurements to separate co-occurring N cycle processes that have counteracting effects on the N isotopes alone. The association of nitrate assimilation with an $^{18}\epsilon_{\text{org}}/^{15}\epsilon_{\text{org}}$ near 1 is based on empirical evidence without full understanding of the physiological mechanisms generating the ratio. A better understanding of the controls on both the magnitude and ratio of N and O isotope effects for nitrate assimilation would strengthen environmental application of N and O isotopes.

Towards this goal, the individual N and O isotope effects for each fractionating step in nitrate assimilation (nitrate reduction, uptake, and efflux) were measured. The reduction of nitrate to nitrite is catalyzed by the intracellular enzyme eukaryotic assimilatory nitrate reductase (eukNR). An N isotope effect of $26.6 \pm 0.2\text{‰}$ and nearly equivalent N and O

fractionation were measured in two distinct forms of eukNR (the NADPH form in cell-free extracts from the fungus *Aspergillus niger* and the NADH form in cell homogenates from the marine diatom *Thalassiosira weissflogii*), suggesting these values will apply to the eukNR family as a whole. These are the first reliable N and O isotope effect measurements for an enzyme that catalyzes the rate-limiting step in nitrate assimilation for all eukaryotic plants and algae.

The N and O isotope effects for nitrate uptake and efflux were measured in the marine diatom *T. weissflogii*. Nitrate uptake and efflux were isolated from nitrate reduction by growing the cells in the presence of tungsten, which substitutes for molybdenum in assimilatory nitrate reductase, yielding an inactive enzyme. The N isotope effects for nitrate uptake and efflux were $2.0 \pm 0.3\%$ and $1.2 \pm 0.4\%$, respectively. The O isotope effect was $2.8 \pm 0.6\%$ for both uptake and efflux, yielding ratios of O to N isotopic fractionation greater than 1 for both processes. In sum, these results confirmed the existing physiological model for isotopic fractionation during nitrate assimilation where the isotope effect associated with intracellular nitrate reductase is high, the isotope effect associated with nitrate uptake is low, and the magnitude of $^{15}\epsilon_{\text{org}}$ depends on the degree to which intracellular fractionation by nitrate reductase is expressed outside the cell by efflux. They also provided the first step in establishing how the $^{18}\epsilon_{\text{org}} \cdot ^{15}\epsilon_{\text{org}}$ near 1 for nitrate assimilation originates at the cellular level.

Finally, the whole cell N and O isotope effects ($^{15}\epsilon_{\text{org}}$ and $^{18}\epsilon_{\text{org}}$) for nitrate assimilation were measured in steady state cultures of *T. weissflogii* to assess how environmental and physiological parameters affected the extent of nitrate efflux and thus the magnitude of $^{15}\epsilon_{\text{org}}$ and $^{18}\epsilon_{\text{org}}$. Steady state cultures ensured $^{15}\epsilon_{\text{org}}$ was not affected by transients in culture conditions and that $^{15}\epsilon_{\text{org}}$ could be related to intracellular nitrate concentration and N and O isotopic composition. As observed in previous studies, $^{15}\epsilon_{\text{org}}$ and intracellular $[\text{NO}_3^-]$ increased under light limitation (a 3-fold and 5-fold increase respectively,

relative to results in non-limited cultures). $^{15}\epsilon_{\text{org}}$ and intracellular $[\text{NO}_3^-]$ were invariant under phosphate limitation. The ratio $^{18}\epsilon_{\text{org}}:^{15}\epsilon_{\text{org}}$ was near 1 under all conditions. In conjunction with previous results from iron- and temperature-limited batch cultures, these results suggest the N and O isotope effects for nitrate assimilation are (i) invariant under most environmental conditions and that (ii) irradiance may be the major driver of variability in $^{15}\epsilon_{\text{org}}$ and $^{18}\epsilon_{\text{org}}$ in the ocean.

The measurements also yielded insight into the regulation of nitrate efflux and suggest a role for intracellular NO_3^- storage in the environment. The near-constant magnitude of $^{15}\epsilon_{\text{org}}$ observed under phosphate-limited growth suggests that the ratio of nitrate efflux to uptake remains 12-15% across a 4-fold increase in growth rate. The constant, non-zero efflux rate this implies suggests a component of efflux is inevitable and correlated with growth rate. The high $^{15}\epsilon_{\text{org}}$ measured under light limitation show that efflux rates increase to 60% of gross nitrate uptake rates, suggesting that an additional component of efflux is correlated with high intracellular $[\text{NO}_3^-]$. The high intracellular $[\text{NO}_3^-]$ observed under light limitation may suggest an adaptation to rapidly changing irradiance in the turbulent mixed layers of the surface ocean: excess intracellular NO_3^- storage may maximize assimilation rates and/or provide an electron sink to dissipate excess energy.

Table of Contents

Abstract.....	i
Table of contents	iv
List of tables	viii
List of figures	ix
Acknowledgements	xii
 Chapter 1.....	 1
1.1. The marine nitrogen cycle	2
1.2. Nitrate N and O Isotopic fractionation and the kinetic isotope effect	3
1.3. The N isotopes of nitrate as tracers of marine N cycle processes	5
1.4. The coupled O and N isotopes of nitrate: isolating the effects of co-occurring N cycle processes	8
1.5. Interpretation of nitrate isotopes: understanding the cellular processes that generate isotopic fractionation	9
1.6. Objectives	12
1.7. References.....	20
Chapter 2:	29
2.1. Introduction.....	31
2.2. Methods	34
2.2.1. Sources of nitrate reductase.....	34
2.2.2. <i>T. weissflogii</i> culture conditions and sampling.....	35
2.2.3. Enzyme assays.....	36
2.2.4. Nitrite determination and removal.....	38
2.2.5. Nitrate isotope analyses	38
2.2.6. Derivation of N and O isotope effects	39

2.3. Results.....	41
2.4. Discussion.....	42
2.4.1. Near equivalent N and O isotope fractionation by eukNR.....	42
2.4.2. The mechanism underlying expression of the N and O isotope effects for nitrate assimilation	44
2.4.3. Support for robust association of nitrate assimilation with an $^{18}\epsilon$: $^{15}\epsilon$ of ~ 1	45
2.4.4. Insights into the catalytic mechanism of eukNR	46
2.4.5. Extension to dissimilatory nitrate reductase NAR.....	51
2.5. References.....	62
Chapter 3.....	70
3.1. Introduction.....	72
3.2. Materials and methods.....	75
3.2.1. Experimental strain.....	75
3.2.2. Culture conditions.....	76
3.2.3. Intracellular nitrate collection.....	77
3.2.4. Nitrate concentration and isotopic composition	78
3.2.5. Particulate nitrogen concentration and isotopic composition.....	81
3.3. Results.....	81
3.3.1. Minimizing nitrate reductase activity with tungstate and growth on NH_4^+	81
3.3.2. Non-zero initial intracellular $[\text{NO}_3^-]$	83
3.3.3. Evolution of intracellular NO_3^-	83
3.3.4. Evolution of intracellular nitrate $\delta^{15}\text{N}$	84
3.3.5. Evolution of intracellular nitrate $\delta^{18}\text{O}$	85
3.4. Discussion.....	85

3.4.1. Fluxes acting on the concentration and N isotopic composition of intracellular nitrate	85
3.4.2. Fluxes acting on the O isotopic composition of intracellular nitrate.....	87
3.4.3. Late increase in intracellular $[\text{NO}_3^-]$	88
3.4.4. Numerical models	88
3.4.5. Analysis of model results.....	98
3.4.6. A $^{15}\epsilon_{\text{in}}$ of 2‰ is consistent with most field and laboratory observations	101
3.4.7. Nitrate uptake makes a significant contribution to organism-level N isotopic fractionation in some conditions	103
3.4.8. Nitrate uptake may cause the O to N ratio for net nitrate assimilation to deviate from the expected ratio of ~ 1 in some conditions	105
3.4.9. Possible origins of the distinct N and O isotopic fractionation associated with nitrate uptake and efflux.....	108
3.4.10. Conclusions.....	110
3.5. References.....	128
Chapter 4:	138
4.1. Introduction.....	140
4.2. Methods	143
4.2.1. Culture conditions.....	143
4.2.2. Sampling.....	145
4.2.3. Dissolved nutrients	146
4.2.4. N and O isotopes of nitrate	146
4.2.5. Particulate organic nitrogen.....	147
4.2.6. Intracellular nitrate.....	148
4.3. Results.....	149

4.3.1. Derivation of N and O isotope effects	149
4.3.2. $^{15}\epsilon_{\text{org}}$	151
4.3.3. Intracellular $[\text{NO}_3^-]$ and $\delta^{15}\text{N}$	152
4.3.4. Fractionation of N and O isotopes of nitrate	152
4.3.5. Cellular quotas and ratios	152
4.3.6. Extracellular $[\text{NO}_3^-]$	153
4.4. Discussion	154
4.4.1. $^{15}\epsilon_{\text{org}}$ is generally invariant	154
4.4.2. Light limitation causes an increase in $^{15}\epsilon_{\text{org}}$ and intracellular $[\text{NO}_3^-]$	157
4.4.3. Relationship between $^{15}\epsilon_{\text{org}}$ and $\delta^{15}\text{N}_{\text{int}}$	159
4.4.4. Nitrate N and O isotopes are fractionated in a ratio near ~ 1	165
4.4.5. Implications for interpretation of nitrate N and O isotope distributions and future work	167
4.5. References	177